



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Register]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for

Display AbstractPlus 20

All: 1

1: [Med Sci Sports Exerc.](#) 1996 Oct;28(10):1327-30.

[Links](#)

Effects of moderate-intensity endurance and high-intensity intermittent training on anaerobic capacity and VO₂max.

[Tabata I](#), [Nishimura K](#), [Kouzaki M](#), [Hirai Y](#), [Ogita F](#), [Miyachi M](#), [Yamamoto K](#).

Department of Physiology and Biomechanics, National Institute of Fitness and Sports, Kagoshima Prefecture, Japan.

This study consists of two training experiments using a mechanically braked cycle ergometer. First, the effect of 6 wk of moderate-intensity endurance training (intensity: 70% of maximal oxygen uptake (VO₂max), 60 min.d⁻¹, 5 d.wk⁻¹) on the anaerobic capacity (the maximal accumulated oxygen deficit) and VO₂max was evaluated. After the training, the anaerobic capacity did not increase significantly ($P > 0.10$), while VO₂max increased from 53 ± 5 ml.kg⁻¹ min⁻¹ to 58 ± 3 ml.kg⁻¹.min⁻¹ ($P < 0.01$) (mean ± SD). Second, to quantify the effect of high-intensity intermittent training on energy release, seven subjects performed an intermittent training exercise 5 d.wk⁻¹ for 6 wk. The exhaustive intermittent training consisted of seven to eight sets of 20-s exercise at an intensity of about 170% of VO₂max with a 10-s rest between each bout. After the training period, VO₂max increased by 7 ml.kg⁻¹.min⁻¹, while the anaerobic capacity increased by 28%. In conclusion, this study showed that moderate-intensity aerobic training that improves the maximal aerobic power does not change anaerobic capacity and that adequate high-intensity intermittent training may improve both anaerobic and aerobic energy supplying systems significantly, probably through imposing intensive stimuli on both systems.

PMID: 8897392 [PubMed - indexed for MEDLINE]

Related Links

- ▶ [Metabolic profile of high intensity intermittent](#) [Med Sci Sports Exerc. 1997]
- ▶ [Ventilatory threshold and VO₂max changes in children following endurance t](#) [Med Sci Sports Exerc. 1989]
- ▶ [Role of thermal factors on aerobic capacity improvements with endurance trainin](#) [J Appl Physiol. 1993]
- ▶ [Effect of training on the anaerobic capacity.](#) [Med Sci Sports Exerc. 1990]
- ▶ [Effectiveness of low-intensity endurance train](#) [Int J Sports Med. 2007]
- ▶ [See all Related Articles...](#)

